

In the Claims:

The following listing reflects amendments to the claims and replaces all prior versions and listings of claims in this application.

1-2. (Cancelled)

²
~~2~~. (Previously presented) The process of claim ¹~~1~~, wherein the protein is a transmembrane protein.

¹
~~1~~. (Currently amended) A process for the preparation of a protein having a molecular weight of about 24kd which specifically binds to the E2 protein of hepatitis C virus, or for the preparation of a functionally equivalent fragment thereof, comprising the steps of:

- i) contacting cells with a preparation of E2;
- ii) obtaining a membrane preparation from cells exhibiting binding to E2; and
- iii) purifying said protein or said functionally equivalent fragment thereof from said preparation.

5-6. (Cancelled)

³
~~3~~. (Currently amended) A process according to any-one either of claims 2-4 ²~~3~~ or ¹~~4~~ wherein the preparation is purified by ammonium sulphate precipitation employing ammonium sulphate at between 33 and 50% saturation.

⁴
~~4~~. (Currently amended) A process according to any-one either of claims 2-4 ²~~3~~ or ¹~~4~~ further comprising at least one hydrophobic interaction chromatography procedure.

⁵
~~5~~. (Currently amended) A process according to any-one either of claims 2-4 ²~~3~~ or ¹~~4~~ further comprising at least one acetone precipitation procedure.

6

~~10~~ (Currently amended) A process for the preparation of a protein having a molecular weight of about 24kd which specifically binds to the E2 protein of hepatitis C virus, or a functionally equivalent fragment thereof, comprising the steps of:

- i) contacting mammalian cells with a preparation of E2;
- ii) obtaining a membrane preparation from the mammalian cells selected for binding to E2;
- iii) precipitating the preparation with ammonium sulphate at less than 33% saturation and retaining the supernatant;
- iv) precipitating the supernatant with ammonium sulphate at between 33 and 50% saturation and retaining the precipitate;
- v) resuspending the precipitate from step iv) in buffer and subjecting the resuspended precipitate to hydrophobic interaction chromatography and recovering the nonretained material to provide said protein having a molecular weight of about 24kd which specifically binds to the E2 protein of hepatitis C virus, or said functionally equivalent fragment thereof.

11-16. (Cancelled)

7

~~17~~ (Currently amended) A diagnostic kit comprising a protein having a molecular weight of about 24 kd, which specifically binds to the E2 protein of hepatitis C virus, or a functionally equivalent fragment thereof, and a labeled HCV E2 protein.

18-20. (Cancelled)

21. (Cancelled)

8

~~22~~ (Previously presented) A method for preparing a protein having a molecular weight of about 24kd which specifically binds to the E2 protein of hepatitis C virus, comprising the steps of:

- i) obtaining a membrane preparation from mammalian cells that bind to E2;
- ii) adding ammonium sulphate to said preparation at less than 33% saturation to produce a precipitate and a supernatant;

iii) adding ammonium sulphate to said supernatant at between 33 and 50% saturation and retaining the precipitate;

iv) resuspending the precipitate from step iii) in buffer and subjecting the resuspended precipitate to hydrophobic interaction chromatography; and

v) recovering said protein.

⁹
~~23~~ (Previously presented) The process of claim ⁸~~22~~ wherein said mammalian cells are MOLT-4 cells.